Supplementary Materials

Selective detection of dopamine with an all PEDOT:PSS Organic Electrochemical Transistor

Isacco Gualandi¹, Domenica Tonelli¹, Federica Mariani¹, Erika Scavetta^{1*}, Marco Marzocchi², Beatrice Fraboni²




Fig. SI 1 Image of the electrochemical (A) and OECT (B) sensors.




Fig SI. 2 CV recorded in 0.1 M PBS, pH 5.5




Fig SI. 3 $\rm I_g\text{-}V_g$ curve recorded in 0.1 M PBS, pH 5.5 . Also the return was record




Fig. SI 4 Response of the sensor in micro molar range




Fig. SI 5. A) Trans-conductance plot obtained for DA for a scan rate equal to 0.002 V s⁻¹. B) Trans-conductance peak values vs dopamine concentration.

Interfering study

In order to gain a better insight into the interfering species effects on the DA response, transconductance curves were recorded at a fixed DA concentration while adding UA and AA to the solution.

In order to minimize the AA and UA contribution to the signal, a high scan rate (0.050 V s⁻¹) was employed. In such a condition, Figure SI 14 A shows the trans-conductance curves recorded in a solution of 0.1 mM DA while UA concentration was increased from 0.1 mM to 6.4 mM. The increase of UA concentration leads to a decrease of DA response probably because UA competes for the occupancy of PEDOT redox sites, causing a saturation of the sensor response. However the DA signal decreases of only 20 % when UA concentration which is 64 times higher than DA concentration.

Figure SI 14 B shows the trans-conductance curves recorded in a solution of 0.1 mM DA while the concentration of AA was increased from 0.1 mM to 0.4 mM. Also in this case the addition of AA produces a decrease of DA signal. The addition of 0.4 mM AA leads to a reduction of dopamine signal of about 20 %. The extent of AA interference is higher than that exerted by UA, probably due to the lower AA oxidation potential.




Fig SI. 6 Trans-conductance curves obtained in a solution containing 0.1 mM DA plus UA (A) and AA (B) at different concentrations.